arXiv:1701.06798v1 [math.RA] 24 Jan 2017

ON KAC'S JORDAN SUPERALGEBRA

ALEJANDRA S. CÓRDOVA-MARTÍNEZ*, ABBAS DAREHGAZANI◇,
AND ALBERTO ELDUQUE*

Abstract. The group-scheme of automorphisms of the ten-dimensional exceptional Kac's Jordan superalgebra is shown to be isomorphic to the semidirect product of the direct product of two copies of $\mathbf{SL}_2$ by the constant group scheme C_2.

This is used to revisit, extend, and simplify, known results on the classification of the twisted forms of this superalgebra and of its gradings.

Kac's ten-dimensional superalgebra K_{10} is an exceptional Jordan superalgebra which appeared for the first time in Kac's classification [Kac77] of the finite dimensional simple Jordan superalgebras over an algebraically closed field of characteristic 0. It was constructed by Lie theoretical terms from a 3-grading of the exceptional simple Lie superalgebra $F(4)$.

A more conceptual definition was given in [BE02] over an arbitrary field $\mathbb{F}$ of characteristic not two (*these assumptions on the ground field will be kept throughout the paper*), using the three-dimensional Kaplansky superalgebra K_3. The even part $(\mathsf{K}_3)_{\bar{0}}$ is a copy of the ground field $\mathbb{F}$: $(\mathsf{K}_3)_{\bar{0}} = \mathbb{F}a$, with $a^2 = a$; while the odd part is a two-dimensional vector space W endowed with a nonzero skew-symmetric bilinear form $(\,|\,)$. The multiplication is determined as follows:

$$a^2 = a, \quad av = \frac{1}{2}v = va, \quad vw = (v|w)a,$$

for any $v, w \in W$. We extend $(\,|\,)$ to a supersymmetric bilinear form on K_3: $(\mathsf{K}_3)_{\bar{0}}$ and $(\mathsf{K}_3)_{\bar{1}}$ are orthogonal, with $(a|a) = \frac{1}{2}$.

Then $\mathsf{K}_{10} = \mathbb{F}1 \oplus \big(\mathsf{K}_3 \otimes_{\mathbb{F}} \mathsf{K}_3\big)$, with $(\mathsf{K}_{10})_{\bar{0}} = \mathbb{F}1 \oplus \big(\mathsf{K}_3 \otimes_{\mathbb{F}} \mathsf{K}_3\big)_{\bar{0}} = \mathbb{F}1 \oplus \mathbb{F}(a \otimes a) \oplus (W \otimes W)$, and $(\mathsf{K}_{10})_{\bar{1}} = \big(\mathsf{K}_3 \otimes_{\mathbb{F}} \mathsf{K}_3\big)_{\bar{1}} = (W \otimes a) \oplus (a \otimes W)$. The multiplication is given by imposing that 1 is the unity, and for homogeneous elements $x, y, z, t \in \mathsf{K}_3$,

$$(x \otimes y)(z \otimes t) = (-1)^{yz}\Big(xz \otimes yt - \frac{3}{4}(x \mid z)(y \mid t)1\Big). \tag{0.1}$$

If the characteristic is 3, then $\mathsf{K}_9 := \mathsf{K}_3 \otimes_{\mathbb{F}} \mathsf{K}_3$ is a simple ideal in K_{10}. Otherwise K_{10} is simple.

It must be remarked that an 'octonionic' construction of K_{10} has been given in [RZ15].

Date: January 23, 2017.

2010 *Mathematics Subject Classification.* Primary 17C40; Secondary 17C70, 17C30.

Key words and phrases. Kac's Jordan superalgebra; group-scheme of automorphisms, twisted forms, gradings.

* Supported by the Spanish Ministerio de Economía y Competitividad—Fondo Europeo de Desarrollo Regional (FEDER) MTM2013-45588-C3-2-P, and by the Diputación General de Aragón—Fondo Social Europeo (Grupo de Investigación de Álgebra). A.S. Córdova-Martínez also acknowledges support from Consejo Nacional de Ciencia y Tecnología (CONACyT, México) through grant 420842/262964.

◇ This research was in part supported by a grant from IPM (No. 94160221) and partially carried out in IPM-Isfahan Branch.

Using the construction above of K_{10} in terms of K_3, the group of automorphisms $\mathsf{Aut}\big(\mathsf{K}_{10}\big)$ was computed in [ELS07]. This was used in [CDM10] to classify gradings on K_{10} over algebraically closed fields of characteristic zero.

Our goal here is to compute the group-scheme of automorphisms of K_{10}, and to use it to revisit, extend, and simplify drastically, the known results on twisted forms and on gradings on K_{10}.

Here we follow the functorial approach (see, for instance, [Wat79]). An affine group-scheme over $\mathbb{F}$ is a representable group-valued functor defined on the category of unital, commutative, associative algebras over $\mathbb{F}$: $\mathrm{Alg}_{\mathbb{F}}$. Thus the group-scheme $\mathbf{Aut}\big(\mathsf{K}_{10}\big)$ is the functor $\mathrm{Alg}_{\mathbb{F}} \to \mathrm{Grp}$ that takes any object R in $\mathrm{Alg}_{\mathbb{F}}$ to the group $\mathsf{Aut}_R\big(\mathsf{K}_{10} \otimes_{\mathbb{F}} R\big)$ (the group of automorphisms of the R-superalgebra $\mathsf{K}_{10} \otimes_{\mathbb{F}} R$, i.e., the group of R-linear isomorphisms preserving the multiplication and the grading), with the natural definition on morphisms.

It turns out that $\mathbf{Aut}\big(\mathsf{K}_{10}\big)$ is isomorphic to a semidirect product $\big(\mathbf{SL}_2 \times \mathbf{SL}_2\big) \rtimes \mathsf{C}_2$ (Theorem 1.1), where C_2 is the constant group scheme attached to the cyclic group of two elements (also denoted by C_2). This extends the result in [ELS07].

Moreover, the canonical projection $\pi : \mathbf{Aut}\big(\mathsf{K}_{10}\big) \to \mathsf{C}_2$ induces an isomorphism of pointed sets $H^1\big(\mathbb{F}, \mathbf{Aut}\big(\mathsf{K}_{10}\big)\big) \to H^1(\mathbb{F}, \mathsf{C}_2)$ (Theorem 2.1). This associates to the isomorphism class of any twisted form of K_{10}, the isomorphism class of a quadratic étale algebra over $\mathbb{F}$, which allows to get easily the classification of twisted forms of K_{10} (Corollary 2.2). Twisted forms of K_{10} were classified, in a completely different way, in [RZ03] and [EO00]. We believe that our approach is more natural.

A simple observation shows that $\big(\mathbf{SL}_2 \times \mathbf{SL}_2\big) \rtimes \mathsf{C}_2$ is also the automorphism group-scheme of $\mathsf{K}_3 \times \mathsf{K}_3$, or of $\mathsf{J}(W) \times \mathsf{J}(W)$, where $\mathsf{J}(W)$ is the Jordan superalgebra of the 'superform' $(\,|\,)$ on $W = W_{\bar{1}}$ above. That is, $\mathsf{J}(W)_{\bar{0}} = \mathbb{F}1$, $\mathsf{J}(W)_{\bar{1}} = W$ and the multiplication is given by

$$1x = x1 = x, \qquad vw = (v \mid w)1,$$

for any $x \in \mathsf{J}(W)$, $v, w \in W$. This observation will be used in the proof of the bijection $H^1\big(\mathbb{F}, \mathbf{Aut}\big(\mathsf{K}_{10}\big)\big) \simeq H^1(\mathbb{F}, \mathsf{C}_2)$.

Finally, given an abelian group G, a G-grading on a superalgebra $\mathcal{A}$ corresponds to a homomorphism of group schemes

$$G^D \longrightarrow \mathbf{Aut}(\mathcal{A}),$$

where G^D is the Cartier dual to the constant group scheme given by G. (See [EK13] for the basic definitions and facts on gradings.)

The classification of G-gradings up to isomorphism on $\mathsf{K}_3 \times \mathsf{K}_3$ (or $\mathsf{J}(W) \times \mathsf{J}(W)$) is an easy exercise (Proposition 3.2), and it follows at once from this the classification of G-gradings, up to isomorphism, on K_{10} (Theorem 3.5), thus extending and simplifying widely the results in [CDM10].

1. The group-scheme of automorphisms

Consider the two-dimensional vector space $W = (\mathsf{K}_3)_{\bar{1}}$, which is endowed with the nonzero skew-symmetric bilinear form $(\,|\,) : W \times W \to \mathbb{F}$. The special linear group-scheme $\mathbf{SL}(W)$ coincides with the symplectic group scheme $\mathbf{Sp}(W)$, which is, up to isomorphism, the group-scheme of automorphisms of K_3 (or of $\mathsf{J}(W)$).

The constant group scheme C_2 acts on $\mathbf{SL}(W) \times \mathbf{SL}(W)$ by swapping the arguments, and hence we get a natural semidirect product $\big(\mathbf{SL}(W) \times \mathbf{SL}(W)\big) \rtimes \mathsf{C}_2$. The group isomorphism Φ in [ELS07, p. 3809] extends naturally to a homomorphism of

affine group-schemes:

$$\begin{aligned}\Phi : \big(\mathbf{SL}(W)\times\mathbf{SL}(W)\big)\rtimes \mathsf{C}_2 &\longrightarrow \mathbf{Aut}\big(\mathsf{K}_{10}\big)\\ (f,g) \quad &\mapsto \quad \Phi_{(f,g)} : \begin{cases} 1 \mapsto 1,\\ a\otimes a\mapsto a\otimes a,\\ v\otimes a\mapsto f(v)\otimes a,\\ a\otimes v\mapsto a\otimes g(v),\\ v\otimes w\mapsto f(v)\otimes g(w),\end{cases}\\ \text{generator of } \mathsf{C}_2 \mapsto \quad &\tau : \begin{cases} 1 \mapsto 1,\\ x\otimes y\mapsto (-1)^{xy} y\otimes x,\end{cases}\end{aligned} \tag{1.1}$$

for any R in $\mathrm{Alg}_{\mathbb{F}}$, $f,g\in\mathbf{SL}(W)(R)$ (i.e., $f,g\in\mathsf{End}_R(W\otimes_{\mathbb{F}}R)\simeq M_2(R)$ of determinant 1), $v,w\in W_R:=W\otimes_{\mathbb{F}}R$, $x,y\in(\mathsf{K}_3)_R$. (Note that a representation of a constant group scheme $\rho:\mathsf{G}\to\mathbf{GL}(V)$ is determined by its behavior over $\mathbb{F}$: $\rho_{\mathbb{F}}:\mathsf{G}\to GL(V)$.)

Theorem 1.1. *Φ is an isomorphism of affine group-schemes.*

Proof. If $\overline{\mathbb{F}}$ denotes an algebraic closure of $\mathbb{F}$, then $\Phi_{\overline{\mathbb{F}}}$ is a group isomorphism [ELS07, Theorem 3.3]. Since $\big(\mathbf{SL}(W)\times\mathbf{SL}(W)\big)\rtimes\mathsf{C}_2\simeq\big(\mathbf{SL}_2\times\mathbf{SL}_2\big)\rtimes\mathsf{C}_2$ is smooth, it is enough to prove that the differential $\mathrm{d}\Phi$ is bijective (see, for instance [EK13, Theorem A.50]). The Lie algebra of $\big(\mathbf{SL}(W)\times\mathbf{SL}(W)\big)\rtimes\mathsf{C}_2$ is $\mathfrak{sl}(W)\times\mathfrak{sl}(W)$, while the Lie algebra of $\mathbf{Aut}\big(\mathsf{K}_{10}\big)$ is the even part of its Lie superalgebra of derivations, which is again, up to isomorphism, $\mathfrak{sl}(W)\times\mathfrak{sl}(W)$ identified naturally with a subalgebra of $\mathsf{End}_{\mathbb{F}}\big(\mathsf{K}_{10}\big)$ (see [BE02, Theorem 2.8]). Moreover, with the natural identifications, $\mathrm{d}\Phi$ is the identity map. □

The last result in this section is the simple observation that $\big(\mathbf{SL}(W)\times\mathbf{SL}(W)\big)\rtimes\mathsf{C}_2$ is also the group-scheme of automorphisms of the Jordan superalgebra $\mathsf{K}_3\times\mathsf{K}_3$ and $\mathsf{J}(W)\times\mathsf{J}(W)$. This will be instrumental in the next section. Its proof is straightforward.

Proposition 1.2. *The natural transformations defined by:*

$$\begin{aligned}\Psi^1 : \big(\mathbf{SL}(W)\times\mathbf{SL}(W)\big)\rtimes \mathsf{C}_2 &\longrightarrow \mathbf{Aut}\big(\mathsf{K}_3\times\mathsf{K}_3\big)\\ (f,g) \quad &\mapsto \quad \Psi^1_{(f,g)} : \begin{cases} (a,0)\mapsto(a,0),\\ (0,a)\mapsto(0,a),\\ (v,w)\mapsto \big(f(v),g(w)\big),\end{cases}\\ \textit{generator of } \mathsf{C}_2 \mapsto \quad &\tau : (x,y)\mapsto(y,x),\end{aligned}$$

for $v,w\in W$ and $x,y\in\mathsf{K}_3$, and

$$\begin{aligned}\Psi^2 : \big(\mathbf{SL}(W)\times\mathbf{SL}(W)\big)\rtimes \mathsf{C}_2 &\longrightarrow \mathbf{Aut}\big(\mathsf{J}(W)\times\mathsf{J}(W)\big)\\ (f,g) \quad &\mapsto \quad \Psi^2_{(f,g)} : \begin{cases} (1,0)\mapsto(1,0),\\ (0,1)\mapsto(0,1),\\ (v,w)\mapsto \big(f(v),g(w)\big),\end{cases}\\ \textit{generator of } \mathsf{C}_2 \mapsto \quad &\tau : (x,y)\mapsto(y,x),\end{aligned}$$

for $v,w\in W$ and $x,y\in\mathsf{J}(W)$, are isomorphisms of group-schemes.

2. Twisted forms

The set of the isomorphism classes of twisted forms of K_{10} is identified with the pointed set $H^1\big(\mathbb{F}, \mathbf{Aut}\big(\mathsf{K}_{10}\big)\big)$. (This is $H^1\big(\overline{\mathbb{F}}/\mathbb{F}, \mathbf{Aut}\big(\mathsf{K}_{10}\big)\big)$ in the notation of [Wat79].)

The key to the classification of twisted forms of K_{10} is the following result:

Theorem 2.1. *The canonical projection* $\pi : \big(\mathbf{SL}(W) \times \mathbf{SL}(W)\big) \rtimes \mathsf{C}_2 \to \mathsf{C}_2$ *induces a bijection of pointed spaces* $\pi_* : H^1\big(\mathbb{F}, \mathbf{Aut}\big(\mathsf{K}_{10}\big)\big) \to H^1(\mathbb{F}, \mathsf{C}_2)$.

Proof. First, the natural section $\iota : \mathsf{C}_2 \to \big(\mathbf{SL}(W) \times \mathbf{SL}(W)\big) \rtimes \mathsf{C}_2$ satisfies $\pi \circ \iota = \mathsf{id}$, so $\pi_* \circ \iota_* = \mathsf{id}$, and π_* is surjective.

Second, the short exact sequence

$$1 \longrightarrow \mathbf{SL}(W) \times \mathbf{SL}(W) \longrightarrow \big(\mathbf{SL}(W) \times \mathbf{SL}(W)\big) \rtimes \mathsf{C}_2 \longrightarrow \mathsf{C}_2 \longrightarrow 1$$

induces an exact sequence (see [Wat79, §18.1]) in cohomology:

$$\begin{aligned} &1 \to \mathrm{SL}(W) \times \mathrm{SL}(W) \to \big(\mathrm{SL}(W) \times \mathrm{SL}(W)\big) \rtimes \mathsf{C}_2 \to \mathsf{C}_2 \\ &\to H^1(\mathbb{F}, \mathbf{SL}(W) \times \mathbf{SL}(W)) \to H^1\big(\mathbb{F}, \big(\mathbf{SL}(W) \times \mathbf{SL}(W)\big) \rtimes \mathsf{C}_2\big) \to H^1(\mathbb{F}, \mathsf{C}_2). \end{aligned}$$

Since $H^1(\mathbb{F}, \mathbf{SL}_2)$ is trivial, this gives an exact sequence

$$1 \longrightarrow H^1\big(\mathbb{F}, \big(\mathbf{SL}(W) \times \mathbf{SL}(W)\big) \rtimes \mathsf{C}_2\big) \to H^1(\mathbb{F}, \mathsf{C}_2),$$

but, in principle, this does not prove that π_* is a bijection. It just says that the only element in $H^1\big(\mathbb{F}, \big(\mathbf{SL}(W) \times \mathbf{SL}(W)\big) \rtimes \mathsf{C}_2\big)$ that is sent by π_* to the trivial element in $H^1(\mathbb{F}, \mathsf{C}_2)$ is the trivial element.

Through the isomorphisms in the previous section, it is enough to prove that π induces a bijection of pointed sets $\pi_* : H^1\big(\mathbb{F}, \mathbf{Aut}\big(\mathsf{J}(W) \times \mathsf{J}(W)\big)\big) \to H^1(\mathbb{F}, \mathsf{C}_2)$.

Write $\mathcal{J} = \mathsf{J}(W) \times \mathsf{J}(W)$ and identify $\big(\mathbf{SL}(W) \times \mathbf{SL}(W)\big) \rtimes \mathsf{C}_2$ with $\mathbf{Aut}(\mathcal{J})$ by means of Ψ^2 in Proposition 1.2. Then $\mathcal{J}_{\bar{0}} = \mathbb{F} \times \mathbb{F}$, and π corresponds to the restriction map $\mathbf{Aut}(\mathcal{J}) \to \mathbf{Aut}(\mathcal{J}_{\bar{0}})$, and the induced map in cohomology $\pi_* : H^1(\mathbb{F}, \mathbf{Aut}(\mathcal{J})) \to H^1(\mathbb{F}, \mathbf{Aut}(\mathcal{J}_{\bar{0}}))$ corresponds to the map that takes the isomorphism class of a twisted form $\mathcal{H}$ of $\mathcal{J}$ to the isomorphism class of $\mathcal{H}_{\bar{0}}$. If $\mathcal{H}_{\bar{0}}$ is trivial ($\mathcal{H}_{\bar{0}} \simeq \mathbb{F} \times \mathbb{F}$) then, by the above, $\mathcal{H}$ is isomorphic to $\mathcal{J}$. Otherwise, $\mathcal{H}_{\bar{0}}$ is a quadratic separable field extension of $\mathbb{F}$, so that $\mathcal{H}_{\bar{1}}$ is a two-dimensional vector space over $\mathcal{H}_{\bar{0}}$, and the multiplication $\mathcal{H}_{\bar{1}} \times \mathcal{H}_{\bar{1}} \to \mathcal{H}_{\bar{0}}$ is $\mathcal{H}_{\bar{0}}$-bilinear and anticommutative, and hence uniquely determined. It turns out that $\mathcal{H}$ is isomorphic to $\mathsf{J}(W) \otimes_{\mathbb{F}} \mathcal{H}_{\bar{0}}$. This proves the injectivity of π_*. □

Corollary 2.2. *Any twisted form of* K_{10} *splits after a quadratic separable field extension, and for any such field extension* $\mathbb{K}$ *of* $\mathbb{F}$*, there is a unique (up to isomorphism) nontrivial twisted form of* K_{10} *which splits over* $\mathbb{K}$.

If $\mathbb{K}$ is a quadratic separable field extension of $\mathbb{F}$, the unique nontrivial twisted form of K_{10} that splits over $\mathbb{K}$ corresponds to the nontrivial element in $H^1(\mathbb{K}/\mathbb{F}, \mathsf{C}_2)$, and hence if the Galois group $\mathrm{Gal}(\mathbb{K}/\mathbb{F})$ is generated by σ ($\sigma^2 = \mathsf{id}$), this twisted form is given by

$$\begin{aligned} \mathsf{K}_{10,\mathbb{K}} &= \{X \in \mathsf{K}_{10} \otimes_{\mathbb{F}} \mathbb{K} : (\tau \otimes \mathsf{id})(X) = (\mathsf{id} \otimes \sigma)(X)\} \\ &= \{X \in \mathsf{K}_{10} \otimes_{\mathbb{F}} \mathbb{K} : (\tau \otimes \sigma)(X) = X\}, \end{aligned} \tag{2.1}$$

where τ is given in (1.1).

Hence, with $\mathbb{K} = \mathbb{F}(\alpha)$, $\alpha^2 \in \mathbb{F}$, $\sigma(\alpha) = -\alpha$, so if $\{u, v\}$ is a symplectic basis of $W = (\mathsf{K}_3)_{\bar{1}}$,

$$\begin{aligned}\big(\mathsf{K}_{10,\mathbb{K}}\big)_{\bar{0}} = \mathbb{F}\text{-span}\Big\langle & 1\otimes 1, (a\otimes a)\otimes 1, (u\otimes u)\otimes\alpha, (v\otimes v)\otimes\alpha, \\ & (u\otimes v - v\otimes u)\otimes 1, (u\otimes v + v\otimes u)\otimes\alpha\Big\rangle, \qquad (2.2)\\ \big(\mathsf{K}_{10,\mathbb{K}}\big)_{\bar{1}} = \mathbb{F}\text{-span}\Big\langle & (a\otimes x + x\otimes a)\otimes 1, (a\otimes x + x\otimes a)\otimes\alpha : x\in\{u,v\}\Big\rangle.\end{aligned}$$

Remark 2.3.

- Both K_3 and $\mathsf{J}(W)$ are rigid: any twisted form of any of these superalgebras is isomorphic to it. This is because $\mathbf{Aut}(\mathsf{K}_3) \simeq \mathbf{Aut}(\mathsf{J}(W)) \simeq \mathbf{SL}(W) \simeq \mathbf{SL}_2$.
- The proof of Theorem 2.1 shows that the twisted forms of $\mathsf{K}_3 \times \mathsf{K}_3$ (respectively $\mathsf{J}(W) \times \mathsf{J}(W)$) are, up to isomorphism, the superalgebras $\mathsf{K}_3 \otimes_{\mathbb{F}} \mathbb{K}$ (resp. $\mathsf{J}(W) \otimes_{\mathbb{F}} \mathbb{K}$), where $\mathbb{K}$ is quadratic étale algebra over $\mathbb{F}$, and any two such forms are isomorphic if and only if so are the corresponding étale algebras.

3. Gradings

Given an abelian group G, a G-grading on a superalgebra $\mathcal{A} = \mathcal{A}_{\bar{0}} \oplus \mathcal{A}_{\bar{1}}$ is a decomposition into a direct sum of subspaces: $\mathcal{A} = \bigoplus_{g\in G} \mathcal{A}_g$, such that $\mathcal{A}_g\mathcal{A}_h \subseteq \mathcal{A}_{gh}$ for any $g, h \in G$, and each homogeneous component is a subspace in the 'super' sense of $\mathcal{A}$: $\mathcal{A}_g = (\mathcal{A}_g \cap \mathcal{A}_{\bar{0}}) \oplus (\mathcal{A}_g \cap \mathcal{A}_{\bar{1}})$. We will write $\deg(x) = g$ in case $0 \neq x \in \mathcal{A}_g$. Two G-gradings $\Gamma : \mathcal{A} = \bigoplus_{g\in G} \mathcal{A}_g$ and $\Gamma' = \bigoplus_{g\in G} \mathcal{A}'_g$, are isomorphic if there is an automorphism $\varphi \in \mathsf{Aut}(\mathcal{A})$ such that $\varphi(\mathcal{A}_g) = \mathcal{A}'_g$ for any $g \in G$.

A grading by G is equivalent to a homomorphism of affine group schemes $G^D \to \mathbf{Aut}(\mathcal{A})$ (see [EK13]), and this shows that two superalgebras with isomorphic group-schemes of automorphisms have equivalent classifications of G-gradings up to isomorphism. Therefore, in order to classify gradings on K_{10} it is enough to classify gradings on $\mathsf{K}_3 \times \mathsf{K}_3$ (or $\mathsf{J}(W) \times \mathsf{J}(W)$), and this is straightforward.

Actually, fix a symplectic basis $\{u, v\}$ of $W = (\mathsf{K}_3)_{\bar{1}}$.

Definition 3.1. Given an abelian group G, consider the following gradings (e denotes the neutral element of G):

- For $g_1, g_2 \in G$, denote by $\Gamma^1(G; g_1, g_2)$ the G-grading given by:
$$\deg(x) = e \text{ for any } x \in \big(\mathsf{K}_3 \times \mathsf{K}_3\big)_{\bar{0}},$$
$$\deg(u, 0) = g_1 = \deg(v, 0)^{-1}, \quad \deg(0, u) = g_2 = \deg(0, v)^{-1}.$$
- For $g, h \in G$ with $h^2 = e \neq h$, denote by $\Gamma^2(G; g, h)$ the G-grading given by:
$$\deg(a, a) = e, \quad \deg(a, -a) = h,$$
$$\deg(u, u) = g = \deg(v, v)^{-1}, \quad \deg(u, -u) = gh = \deg(v, -v)^{-1}.$$

Proposition 3.2. *Any grading on $\mathsf{K}_3 \times \mathsf{K}_3$ by the abelian group G is isomorphic to either $\Gamma^1(G; g_1, g_2)$ or to $\Gamma^2(G; g, h)$ (for some g_1, g_2 or g, h in G).*

Moreover, no grading of the first type ($\Gamma^1(G; g_1, g_2)$) is isomorphic to a grading of the second type ($\Gamma^2(G; g, h)$), and

- *$\Gamma^1(G, g_1, g_2)$ is isomorphic to $\Gamma^1(G; g'_1, g'_2)$ if and only if the sets $\{g_1, g_1^{-1}, g_2, g_2^{-1}\}$ and $\{g'_1, (g'_1)^{-1}, g'_2, (g'_2)^{-1}\}$ coincide.*

- $\Gamma^2(G;g,h)$ *is isomorphic to* $\Gamma^2(G;g',h')$ *if and only if* $h'=h$ *and* $g'\in\{g,gh,g^{-1},g^{-1}h\}$.

Proof. Any G-grading on $\mathcal{J}:=\mathsf{K}_3\times\mathsf{K}_3$ induces a G-grading on $\mathcal{J}_{\bar{0}}$, which is isomorphic to $\mathbb{F}\times\mathbb{F}$, and hence we are left with two cases:

(1) The grading on $\mathcal{J}_{\bar{0}}$ is trivial, i.e., $\mathcal{J}_{\bar{0}}$ is contained in the homogeneous component $\mathcal{J}_e$. Then, with $W=(\mathsf{K}_3)_{\bar{1}}$, both $W\times 0=(a,0)\mathcal{J}_{\bar{1}}$ and $0\times W=(0,a)\mathcal{J}_{\bar{1}}$ are graded subspaces of $\mathcal{J}_{\bar{1}}$. Hence we can take bases $\{u_i,v_i\}$ of W, $i=1,2$, such that $\{(u_1,0),(v_1,0),(0,u_2),(0,v_2)\}$ is a basis of $\mathcal{J}_{\bar{1}}$ consisting of homogeneous elements. We can adjust v_i, $i=1,2$, so that $(u_i\mid v_i)=1$. If $\deg(u_i,0)=g_i$, $i=1,2$, the grading is isomorphic to $\Gamma^1(G;g_1,g_2)$.

(2) The grading on $\mathcal{J}_{\bar{0}}$ is not trivial. Then there is an element $h\in G$ of order 2 such that $\deg(a,a)=e$ and $\deg(a,-a)=h$. (Note that (a,a) is the unity element of $\mathcal{J}_{\bar{0}}$ ($\simeq\mathbb{F}\times\mathbb{F}$), so it is always homogeneous of degree e.) As $\mathcal{J}_{\bar{0}}=(\mathcal{J}_{\bar{1}})^2$, there are homogeneous elements $(u_1,u_2),(v_1,v_2)\in\mathcal{J}_{\bar{1}}$ such that $(u_1,u_2)(v_1,v_2)=(a,a)$. If $g=\deg(u_1,u_2)$, this grading is isomorphic to $\Gamma^2(G;g,h)$.

The conditions for isomorphism are clear. □

Any grading $\Gamma^1(G;g_1,g_2)$ is a coarsening of the grading $\Gamma^1\big(\mathbb{Z}^2;(1,0),(0,1)\big)$, while any grading $\Gamma^2(G;g,h)$ is a coarsening of the grading $\Gamma^2\big(\mathbb{Z}\times\mathbb{Z}_2;(1,\bar{0}),(0,\bar{1})\big)$, where $\mathbb{Z}_2=\mathbb{Z}/2\mathbb{Z}$. As an immediate consequence, we obtain the next result.

Corollary 3.3. *Up to equivalence, there are exactly two fine gradings on* $\mathsf{K}_3\times\mathsf{K}_3$*:* $\Gamma^1\big(\mathbb{Z}^2;(1,0),(0,1)\big)$ *and* $\Gamma^2\big(\mathbb{Z}\times\mathbb{Z}_2;(1,\bar{0}),(0,\bar{1})\big)$.

Proposition 3.2 and Corollary 3.3 have completely similar counterparts for $\mathsf{J}(W)\times\mathsf{J}(W)$.

In order to transfer these results to Kac's superalgebra K_{10}, take into account that K_{10} is generated by its odd part, as $\big((\mathsf{K}_{10})_{\bar{1}}\big)^2=(\mathsf{K}_{10})_{\bar{0}}$, so any grading is determined by its restriction to the odd part, and use the commutativity of the diagram

$$\begin{array}{ccccc}
\mathbf{Aut}\big(\mathsf{K}_3\times\mathsf{K}_3\big) & \xleftarrow{\Psi^1} & \big(\mathbf{SL}(W)\times\mathbf{SL}(W)\big)\rtimes\mathsf{C}_2 & \xrightarrow{\Phi} & \mathbf{Aut}\big(\mathsf{K}_{10}\big)\\
\downarrow & & & & \downarrow\\
\mathbf{GL}(W\times W) & & \xrightarrow{\qquad\simeq\qquad} & & \mathbf{GL}\big((W\otimes a)\oplus(a\otimes W)\big)
\end{array}$$

where the vertical arrows are given by the restrictions to the odd parts, and the bottom isomorphism is given by the natural identification $W\times W\to(W\otimes a)\oplus(a\otimes W)$, $(v,w)\mapsto v\otimes a+a\otimes w$.

Thus Definition 3.1 transfers to K_{10} as follows:

Definition 3.4. Given an abelian group G, consider the following G-gradings on K_{10}:

- For $g_1,g_2\in G$, denote by $\Gamma^1_{\mathsf{K}_{10}}(G;g_1,g_2)$ the G-grading determined by:
$$\deg(u\otimes a)=g_1=\deg(v\otimes a)^{-1},\quad \deg(a\otimes u)=g_2=\deg(a\otimes v)^{-1}.$$
- For $g,h\in G$ with $h^2=e\neq h$, denote by $\Gamma^2_{\mathsf{K}_{10}}(G;g,h)$ the G-grading determined by:
$$\begin{aligned}\deg(u\otimes a+a\otimes u)&=g=\deg(v\otimes a+a\otimes v)^{-1},\\ \deg(u\otimes a-a\otimes u)&=gh=\deg(v\otimes a-a\otimes v)^{-1}.\end{aligned}$$

And Proposition 3.2 and Corollary 3.3 are now transferred easily to K_{10}:

Theorem 3.5. *Any grading on* K_{10} *by the abelian group* G *is isomorphic to either* $\Gamma^1_{\mathsf{K}_{10}}(G; g_1, g_2)$ *or to* $\Gamma^2_{\mathsf{K}_{10}}(G; g, h)$ *(for some* g_1, g_2 *or* g, h *in* G*).*

No grading of the first type is isomorphic to a grading of the second type, and

- $\Gamma^1_{\mathsf{K}_{10}}(G; g_1, g_2)$ *is isomorphic to* $\Gamma^1_{\mathsf{K}_{10}}(G; g_1', g_2')$ *if and only if the sets* $\{g_1, g_1^{-1}, g_2, g_2^{-1}\}$ *and* $\{g_1', (g_1')^{-1}, g_2', (g_2')^{-1}\}$ *coincide.*
- $\Gamma^2_{\mathsf{K}_{10}}(G; g, h)$ *is isomorphic to* $\Gamma^2_{\mathsf{K}_{10}}(G; g', h')$ *if and only if* $h' = h$ *and* $g' \in \{g, gh, g^{-1}, g^{-1}h\}$.

Moreover, there are exactly two fine gradings on K_{10} *up to equivalence, namely* $\Gamma^1_{\mathsf{K}_{10}}\big(\mathbb{Z}^2; (1,0), (0,1)\big)$ *and* $\Gamma^2_{\mathsf{K}_{10}}\big(\mathbb{Z} \times \mathbb{Z}_2; (1, \bar{0}), (0, \bar{1})\big)$.

Remark 3.6. Any grading on K_{10} by an abelian group G extends naturally to a grading by either $\mathbb{Z} \times G$ or $\mathbb{Z}_2^2 \times G$ on the exceptional simple Lie superalgebra $F(4)$, which is obtained from K_{10} using the well-known Tits-Kantor-Koecher construction. However, not all gradings on $F(4)$ are obtained in this way.

References

[BE02] G. Benkart and A. Elduque, *A new construction of the Kac Jordan superalgebra*, Proc. Amer. Math. Soc. **130** (2002), no. 11, 3209–3217.

[CDM10] A.J. Calderón Martín, C. Draper Fontanals, and C. Martín González, *Gradings on the Kac superalgebra*, J. Algebra **324** (2010), no. 12, 3249–3261.

[EK13] A. Elduque and M. Kochetov, *Gradings on simple Lie algebras*, Mathematical Surveys and Monographs **189**, American Mathematical Society, Providence, RI, 2013.

[EO00] A. Elduque and S. Okubo, *Pseudo-composition superalgebras*, J. Algebra **227** (2000), no. 1, 1–25.

[ELS07] A. Elduque, J. Laliena, and S. Sacristán, *The Kac Jordan superalgebra: automorphisms and maximal subalgebras* Proc. Amer. Math. Soc. **135** (2007), no. 12, 3805–3813.

[Kac77] V.G. Kac, *Classification of simple* $\mathbb{Z}$*-graded Lie superalgebras and simple Jordan superalgebras*, Comm. Algebra **5** (1977), no. 13, 1375–1400.

[RZ03] M.L. Racine and E.I. Zelmanov, *Simple Jordan superalgebras with semisimple even part*, J. Algebra **270** (2003), no. 2, 374–444.

[RZ15] M.L. Racine and E.I. Zelmanov, *An octonionic construction of the Kac superalgebra K10*, Proc. Amer. Math. Soc. **143** (2015), no. 3, 1075–1083.

[Wat79] W.C. Waterhouse, *Introduction to affine group schemes*, Graduate Texts in Mathematics **66**. Springer-Verlag, New York-Berlin, 1979.

Departamento de Matemáticas e Instituto Universitario de Matemáticas y Aplicaciones, Universidad de Zaragoza, 50009 Zaragoza, Spain

E-mail address: sarina.cordova@gmail.com

Department of Mathematics, University of Isfahan, Isfahan, Iran, P.O.Box: 81746-73441 and School of Mathematics, Institute for Research in Fundamental Sciences (IPM), P.O. Box: 19395-5746

E-mail address: abbas.darehgazani@gmail.com

Departamento de Matemáticas e Instituto Universitario de Matemáticas y Aplicaciones, Universidad de Zaragoza, 50009 Zaragoza, Spain

E-mail address: elduque@unizar.es